

## Scottish Government Central Analysis Division

### Coronavirus (Covid-19): modelling the epidemic in Scotland (Issue No. 107)

#### Background

This is a report on the Scottish Government modelling of the spread and level of Covid-19 in Scotland. This updates the previous publication on modelling of Covid-19 in Scotland published on 18th August 2022. The estimates in this document provide an overview of the situation regarding the virus and help the Scottish Government, the health service and the wider public sector plan ahead.

#### Key Points

- The reproduction rate  $R$  in Scotland is currently estimated as being between 0.8 and 1.0, as at 16th August. The lower limit has increased since the last publication while the upper limit is unchanged.
- The daily growth rate for Scotland is currently estimated as between -4% and 0% as at 16th August. Both the lower limit and the upper limit have increased since the last publication.
- The number of new daily infections for Scotland is estimated as being between 90 and 304 per 100,000 people as at 16th August.
- Average contacts from the most recent wave of the Scottish Contact Survey (18th August - 24th August) indicate an average of 5.7 contacts. This has increased by 13% compared to the previous wave of the survey (4th August - 10th August) where average contacts were 5.1. Prior to the Covid-19 pandemic, the average daily contacts for adults in the UK were reported to be 10.8.
- Mean contacts in the work setting have decreased by 20% over the last two weeks whilst mean contacts within the home have increased by 6%. Mean contacts in the other setting (contacts outside home, school and work) have remained at a similar level over the same

period. Average contacts have increased in all age groups in the 18-59 range, with contacts in the 18-29 age group increasing by 128%.

- Mean contacts have increased by 38% in the 18-29 age group and by 16% in those aged 70 and over. Mean contacts have remained at a similar level in all other age groups.
- The highest average number of interactions are observed between the 18-29 age group and those under the age of 18.
- The largest decrease in the proportion of participants visiting different locations is seen in those visiting non-essential shops. This decreased from 55% to 52%. The largest increase was seen in proportion of participants using public transport, increasing from 32% to 36%.
- The percentage of people wearing a face covering where they have at least one contact outside of the home has decreased from 26% to 25% since the last wave of the survey.
- 30% of people had taken at least one lateral flow test in the previous 7 days, decreasing from 31% in the previous wave of the survey pertaining to the 4th August - 10th August.
- The SPI-M-O consensus view is that by 11th October, daily hospitalisations from Covid-19 in Scotland are estimated to be between 3 and 122, and hospital occupancy is estimated to be between 110 and 1142.
- Analysis by the Edinburgh University Roslin Institute indicates that there remains a substantial variation in geographical risk of infection when considering the average recorded census age and booster vaccination uptake.
- The data show a continued decline in reported tests across all deprivation deciles, as well as a decline in reported positive tests and overall positivity. There is no clear trend in positivity with respect to deprivation. The volume of PCR testing remains very low.
- The distribution of lateral flow/LFD tests being reported continues to vary substantially by both age and deprivation status, with many fewer tests reported in younger adults across all deciles of deprivation, and for children in more deprived deciles (with some deciles now reporting zero negative lateral flow tests over the course of a week). The high level of LFD positivity in the latter category is marked, and when compared to the high number of positives amongst the least deprived, consistent with under ascertainment

being concentrated (and in turn higher circulation than the cases data suggest) in particular groups.

- Nationwide, during the period 12th – 25th August, Wastewater Covid-19 levels remained relatively stable in the range of 55 to 73 million gene copies per person per day (Mgc/p/d), similar to the previous two-week period (28th July – 11th August).

## **Overview of Scottish Government Modelling**


Modelling outputs are provided here on the current epidemic in Scotland as a whole, based on a range of methods. Because it takes a little over three weeks on average for a person who catches SARS-CoV-2 (the causative agent of Covid-19) to show symptoms, become sick and either die or recover, there is a time lag in what our model can tell us about any changes in the epidemic.

The Scottish Government presents its modelling outputs to the Epidemiology Modelling Review Group (EMRG). These outputs are included (shown in green) in both Figure 1 and Figure 2. Outputs from other modellers are shown in either cyan or purple. The UKHSA consensus range is the rightmost (in red).

The R value and growth rates are also estimated by several independent modelling groups based in universities and the UKHSA. Estimates are considered, discussed and combined at the EMRG, which sits within UKHSA. These are based on data up to 29th August.

The consensus view of the UKHSA across these methods was that the value of R in Scotland is between 0.8 and 1.0 as of 16th August 2022 (Figure 1). The lower limit has increased since the previous publication while the upper limit is unchanged. R is an indicator that lags by two to three weeks.

Figure 1: Estimates of  $R_t$  for Scotland as of 16th August, including 90% confidence intervals, produced by EMRG




Source: EMRG

The consensus from UKHSA is that the growth rate in Scotland is between -4% and 0% per day as at 16th August. Both the upper and lower limits have increased since the last publication.

The various groups which report to EMRG use different sources of data in their models to produce estimates of incidence (Figure 2). The consensus view of the UKHSA across these methods was that the incidence of new daily infections in Scotland was between 90 and 304 new infections per 100,000 as of 16th August. This equates to between 4,900 and 16,600 people becoming infected each day in Scotland.

Figure 2: Estimates of incidence for Scotland, as at 16th August, including 90% confidence intervals, produced by EMRG



Source: EMRG

### What we know about how people's contact patterns have changed

From this week's publication, Scottish Contact Survey contact matrices are published as [open data](#). The dataset contains mean contacts between age groups for all waves of the survey since its inception in August 2020 and will be updated fortnightly to add data for the latest survey wave.

Average contacts from the most recent wave of the Scottish Contact Survey (18th August - 24th August) indicate an average of 5.7 contacts. This has increased by 13% compared to the previous wave of the survey (4th August - 10th August) where average contacts were 5.1, as seen in Figure 3.

Prior to the Covid-19 pandemic, the average daily contacts for adults in the UK were reported to be 10.8 from the UK-wide POLYMOD study<sup>1</sup>.

Mean contacts in the work setting have decreased by 20% over the last two weeks whilst mean contacts within the home have increased by 6%. Mean contacts in the other setting (contacts outside home, school and work) have remained at a similar level over the same period.

<sup>1</sup> <https://bmcmmedicine.biomedcentral.com/track/pdf/10.1186/s12916-020-01597-8>

Figure 3: Mean contacts per day for adults in Scotland (truncated to 100 contacts per participant)






Figure 4 shows how contacts change across age group and setting. Mean contacts have increased by 38% in the 18-29 age group and by 16% in those aged 70 and over. Mean contacts have remained at a similar level in all other age groups.

Figure 4: Mean contacts per day for adults in Scotland (truncated to 100 contacts per participant) by age group and setting



The heatmaps in Figure 5 show the mean overall contacts between age groups for the surveys relating to 4th August - 10th August and 18th August - 24th August and the difference between these periods. The highest average number of interactions are observed between the 18-29 age group and those under the age of 18.

Figure 5: Mean contacts per day between age groups (truncated to 100 contacts per participant)



The largest decrease in the proportion of participants visiting different locations is seen in those visiting non-essential shops. This decreased from 55% to 52%. The largest increase was seen in proportion of participants using public transport, increasing from 32% to 36%.

**Figure 6: Locations visited by participants at least once**





Figure 7 shows the percentage of people wearing a face covering where they have at least one contact outside of the home. This has decreased from 26% to 25% since the last wave of the survey.

**Figure 7: Percentage of participants wearing a face covering (with at least one contact outside of the home)**




In the survey pertaining to 18th August - 24th August, 30% of people had taken at least one lateral flow test in the previous 7 days,



decreasing from 31% in the previous wave of the survey pertaining to the 4th August - 10th August.

*Figure 8: Percentage of participants who had taken at least one lateral flow test in the last 7 days*



### **What the modelling tells us about projections of hospitalisations and hospital occupancy in the medium term**

SPI-M-O produces projections of the epidemic (Figure 9 and Figure 10), combining estimates from several independent models. These projections are not forecasts or predictions. They represent a scenario in which the trajectory of the epidemic continues to follow the trends that were seen in the data available to 29th August and do not include the effects of any future policy or behavioural changes. The delay between infection, developing symptoms and the need for hospital care means they cannot fully reflect the impact of behaviour changes in the two to three weeks prior to 29th August. The projections include the potential impact of vaccinations over the next few weeks. Modelling groups have used their expert judgement and evidence from UKHSA, Scottish

Universities & Public Health Scotland, and other published efficacy studies when making assumptions about vaccine effectiveness

Figure 9: SPI-M-O medium-term projection of daily hospitalisations in Scotland, at 50% and 90% credible intervals





Figure 10 shows the SPI-M-O consensus on hospital occupancy. Hospital occupancy is determined by the combination of admissions and length of stay, the latter of which is difficult to model with confidence.

Figure 10: SPI-M-O medium-term projection of hospital occupancy in Scotland, at 50% and 90% credible intervals



The SPI-M-O consensus view is that by 11th October, daily hospitalisations from Covid-19 in Scotland are estimated to be between

3 and 122, and hospital occupancy is estimated to be between 110 and 1142.

### **Summary of spatial analysis of Covid-19 spread in Scotland**

Researchers at the Edinburgh Roslin Institute have conducted spatial analysis of COVID-19 Spread in Scotland. A summary of findings from data up to 25th August 2022 is included here.

Rates of LFD and PCR testing are stabilising, albeit at very low levels (1-2 LFD tests reported, per week, per 1,000).

The distribution of lateral flow/LFD tests being reported varies substantially by both age and deprivation status, with many fewer tests reported in younger adults across all deciles of deprivation, and for children in more deprived deciles. The high level of LFD positivity in the latter category is marked, and when compared to the high number of positives amongst the least deprived, suggests that ascertainment may be lower in younger people in deprived areas.

Figure 11: Variation in testing outcomes comparing Lateral Flow and PCR testing considering age and deprivation status of the data zone of record based on data in week ending 20th August 2022

Testing and positivity - week ending 2022-08-20





Figure 12: Variation in testing outcomes comparing Lateral Flow and PCR testing, separated by deprivation



## What can analysis of wastewater samples tell us about local outbreaks of Covid-19 infection?

Levels of Covid-19 RNA in wastewater (WW) collected at a number of sites around Scotland are adjusted for population and local changes in intake flow rate (or ammonia levels where flow is not available). See Technical Annex in Issue 34 of Scottish Government Research Findings for the methodology. These reports are based on the most recent data available. Future updates to data may lead to small retrospective changes.


Nationwide, during the period 12th – 25th August, Wastewater Covid-19 levels remained relatively stable in the range of 55 to 73 million gene copies per person per day (Mgc/p/d), similar to the previous two-week period (28th July – 11th August).

In Figure 13, we show wastewater RNA levels up to 25th August 2022, with the ONS Coronavirus Infection Survey (CIS) estimates up to 16th August 2022 superimposed. The CIS data indicate a continuing

decrease in the percentage of population testing positive for COVID-19 up to 16th August.

Site level data also suggest that Wastewater Covid-19 levels in the Edinburgh area may be increasing over the past two weeks, unlike other major cities.

Figure 13: National running average trends in wastewater Covid-19 as of 25th August<sup>2</sup>



### Looking to the future

What may happen in the future around SARS-CoV-2 is uncertain and therefore there are a number of possible Covid-19 futures that may occur in the future. For example, the current Omicron wave may dissipate leaving low levels of Covid-19, or a new variant may emerge potentially having vaccine escape or increased severity, or people's behaviours may change. One approach to this uncertainty is to model alternative versions of the future through the development of different Covid-19 scenarios.

Given what we know about Covid-19 these possible futures range from a world where immunity reduces Covid-19 hospitalisations and ICU to low levels, through to variant world where a variant with immune escape

<sup>2</sup> For this graph, a wastewater RNA average using the last 7 days of data is computed at every sampling date. Prevalence estimates and 95% confidence intervals from the ONS Coronavirus Infection Survey is overlaid, with a scale chosen to approximately match Jan/Feb trends in WW Covid-19.

enters Scotland and Covid-19 hospitalisations and ICU could increase. In between these two extremes there could be possible futures where people's behaviour becomes polarised between those who continue with Covid-19 precautions e.g. hand washing etc. and those who do not.

The scenarios we provide in the next section look at what could happen for planning purposes, not to forecast what will happen. The assumptions are based on our most up to date knowledge, but do not include the effect of future changes in treatment of Covid-19 e.g. widespread use of antivirals or changes in behaviour in response to high levels of infections e.g. in variant world. Therefore, in the most extreme scenarios the peak may be lower than suggested if behaviour or restrictions changed.


There is no linear progression between the worlds and all are plausible. Each world inherently contains a different threat level requiring a different approach to management.

## Immune World

In this possible future vaccines and natural immunity are effective at keeping Covid-19 at low levels. New variants may emerge in Scotland but for the foreseeable future infections are based around Omicron.

Infections may decrease from current levels over the coming weeks and months to very low levels. Likewise hospital and ICU occupancy may follow this trend relieving the pressure on healthcare services. Issues with new variants are not considered in this world and therefore levels of infections remain low.

*Figure 14: Potential infections, hospital occupancy and ICU occupancy trajectory in Immune World*



In Immune world Covid-19 in Scotland reduces below epidemic levels, becoming endemic. Cases of Covid-19 therefore spring up only as rare outbreaks which are controlled through public health measures. People's lives

return to something close to normality e.g. physical distancing is not needed but people still choose to self-isolate and hygiene is good. As vaccines are effective, take-up of first/second/third doses are good and boosters become part of an annual cycle like flu. The numbers of people who need medical treatment or hospitalisation for Covid-19 remain low.


The focus moves away from Covid-19 response and into recovery. This includes addressing learning losses, treating Long Covid and working through the hospital backlog. Wellbeing measures improve with reduced anxiety and increased happiness. Those from the highest risk groups feel they can reintegrate without government interventions. The economy continues to recover from the effects of Covid-19. Travellers do not face significant issues with trips overseas.

## Polarised world

In this world, vaccines and natural immunity are effective at reducing infections. The approach followed relies on individual risk assessment and behaviours. However, society becomes polarised as some continue to take up vaccines and exercise caution while others are more reluctant. Covid-19 becomes a disease associated with those who do not or cannot get full vaccine benefit and do not or cannot adopt a risk based approach maintaining baseline measures.

Impacts on hospital/ICU occupancy are uncertain but levels may be higher than has been observed in summer 2020 and 2021 and higher than what may happen in Immune world (see Figure 15).

*Figure 15: Potential infections, hospital occupancy and ICU occupancy trajectory in Polarised World*



Cases of Covid-19 spring up and are hard to control in those who are not vaccinated or vulnerable. People's lives return to a "new normal" but, due to polarised groups in society with some following and some not following guidance, infections remain.



Vaccines are effective so older and more vulnerable people come forward for future doses in high numbers.

The focus remains on Covid-19 and the shift onto recovery is slower. Existing learning losses are harder to rectify and continue to accrue due to infections within education settings. The hospital backlog is difficult to address as hospitals are still dealing with Covid-19 cases. The population becomes polarised in to those whose wellbeing improves e.g. lower risk people and those whose wellbeing deteriorates e.g. higher risk or poorer people whose levels of anxiety increase as Covid-19 circulates. They continue to experience greater illness, greater poverty or disruption to their income. The economy continues to be impacted from the effects of Covid-19.


### **Variant world – vaccine escape with same severity as Delta**

In this possible future a variant with vaccine escape emerges in Scotland presenting a challenge even for fully vaccinated people. This new variant leads to increased transmission, but not to increased severity compared to previous variants. In this scenario other NPIs may need to be put in place for a short time. This world is similar to what has happened in Scotland with the emergence of Omicron.

Omicron may be reduced to low levels within Scotland as a new variant takes over. This causes a new wave of Covid-19 infections as well as increases in hospital and ICU occupancy. People's lives are disrupted due to the increasingly high levels of infections leading to time off work ill or isolating.

To show the potential impact assume a new variant appears in Scotland in the October holidays as people return to work and school. The timing is uncertain and a potential new variant may appear sooner than the summer or significantly later but has currently been lined up with the summer holidays to show illustratively what could happen. The new variant may cause Omicron infections to decrease significantly or disappear entirely (and is not shown). The new variant is modelled with similar transmissibility and vaccine escape as Omicron with severity characteristics similar to Delta. It could lead to high levels of infections leading to hospital occupancy rising above capacity restrictions. With sustained high levels of infection we could again see increased staff absences in a number of sectors that were affected by this in the recent Omicron wave.

*Figure 16: Potential infections, hospital occupancy and ICU occupancy trajectory in Variant World with the same severity as Delta*



The focus remains on Covid-19 and it is hard to shift on to recovery. Continued infections within education settings and staff shortages may impact schools. The Covid-19 strain on hospitals is high due to the very high numbers of infections and workforce pressures grow making it difficult to address the hospital backlog. Wellbeing measures deteriorate with people reporting low happiness and general 'tiredness with it all'. The economy continues to be impacted from the effects of Covid-19 with many people off work. Travellers may not want to come to the UK as the new variant sweeps through.


### **Variant world – vaccine escape with increased severity compared to Delta**

As with the other example of Variant world, a new variant appears in Scotland in the October holidays as people return to work and school. The timing is uncertain but has currently been lined up with the summer holidays where reduced travel restrictions may make it more likely that a new variant is brought into Scotland.

The new variant may cause Omicron infections to decrease significantly or disappear entirely (and this is not shown on the graph). It is modelled with similar transmissibility and vaccine escape as Omicron with severity characteristics 50% higher than Delta, purely for illustrative purposes.

It could lead to high levels of infections leading to hospital occupancy rising significantly. With sustained high levels of infection we could again see increased staff absences in a number of sectors that were affected by this in the recent Omicron wave.

*Figure 17: Potential infections, hospital occupancy and ICU occupancy trajectory in Variant World with the increased severity compared to Delta*



The focus remains on Covid-19 and it is hard to shift on to recovery. Continued infections within education settings and staff shortages may impact schools. The Covid-19 strain on hospitals is high due to the very high numbers of infections and workforce pressures grow making it difficult to address the hospital backlog. Wellbeing measures deteriorate with people reporting low happiness and general 'tiredness with it all'. The economy continues to be impacted from the effects of Covid-19 with many people off work. Travellers may not want to come to the UK as the new variant sweeps through.

### **What next?**

Archiving of models is currently being undertaken via the [Data Science Scotland](#) GitHub organisation. Details of the [Epidemia](#) model has most recently been made available. More models will be added over the coming weeks - see the Technical Annex of issue 96 for further details.

## Technical Annex

### What levels of Covid-19 are indicated by wastewater data?

Table 1 provides population weighted daily averages for normalised wastewater Covid-19 levels in the weeks ending 18th and 25th August 2022<sup>3</sup>, with no estimate for error. This is given in Million gene copies per person per day. Coverage is given as percentage of inhabitants in each local authority covered by a wastewater Covid-19 sampling site delivering data during this period.

*Table 1. Average Covid-19 wastewater levels (Mgc/p/d)<sup>4</sup>.*

Local Authority (LA)	w/e 18th August	w/e 25th August	Coverage
Aberdeen City	69	73	99 %
Aberdeenshire	50	48	50 %
Angus	46	74	68 %
Argyll and Bute	14	156	23 %
City of Edinburgh	88	108	98 %
Clackmannanshire	39	67	92 %
Dumfries and Galloway	20	43	39 %
Dundee City	52	81	100 %
East Ayrshire	60	100	72 %
East Dunbartonshire	57	84	99 %
East Lothian	71	89	74 %
East Renfrewshire	22	75	89 %
Falkirk	40	68	96 %
Fife	66	58	84 %
Glasgow City	34	67	98 %
Highland	45	61	48 %
Inverclyde	34	36	98 %
Midlothian	82	107	88 %
Moray	96	67	70 %
Na h-Eileanan Siar	–	–	0 %
North Ayrshire	39	59	92 %
North Lanarkshire	50	44	95 %
Orkney Islands	–	112	34 %
Perth and Kinross	40	37	45 %
Renfrewshire	55	51	97 %
Scottish Borders	28	30	59 %
Shetland Islands	–	–	0 %
South Ayrshire	59	103	77 %
South Lanarkshire	41	44	88 %

<sup>3</sup> Results from the most recent samples taken on Friday 26th August were not available due to technical difficulties and hence our results are not as up to date as usual

<sup>4</sup> Coverage as for week ending 25th August 2022.

Stirling	37	18	63 %
West Dunbartonshire	53	108	98 %
West Lothian	33	42	95 %

This publication will be available in accessible HTML on the [gov.scot](http://www.gov.scot) website

© Crown copyright 2022

You may re-use this information (excluding logos and images) free of charge in any format or medium, under the terms of the Open Government Licence. To view this licence, visit <http://www.nationalarchives.gov.uk/doc/open-government-licence/> or e-mail: [psi@nationalarchives.gsi.gov.uk](mailto:psi@nationalarchives.gsi.gov.uk). Where we have identified any third party copyright information you will need to obtain permission from the copyright holders concerned.

The views expressed in this report are those of the researchers and do not necessarily represent those of the Scottish Government or Scottish Ministers.

This document is also available from our website at [www.gov.scot](http://www.gov.scot).  
ISBN: 978-1-80435-910-5

The Scottish Government  
St Andrew's House  
Edinburgh  
EH1 3DG

Produced for  
the Scottish Government  
by APS Group Scotland  
PPDAS1151702 (09/22)  
Published by  
the Scottish Government,  
September 2022



ISBN 978-1-80435-910-5

Web Publication

PPDAS1151702 (09/22)