



Scottish Government Central Analysis Division

Coronavirus (Covid-19): modelling the epidemic in Scotland (Issue No. 111)

Background

This is a report on the Scottish Government modelling of the spread and level of Covid-19 in Scotland. This updates the previous publication on modelling of Covid-19 in Scotland published on 13th October 2022. The estimates in this document provide an overview of the situation regarding the virus and help the Scottish Government, the health service and the wider public sector plan ahead.

Key Points

- The reproduction rate R in Scotland is currently estimated as being between 0.8 and 1.1, as at 11th October. Both the lower limit and the upper limit have decreased since the last publication.
- The daily growth rate for Scotland is currently estimated as between -4% and 3% as at 11th October. The lower limit has decreased since the last publication. The upper limit is unchanged.
- The number of new daily infections for Scotland is estimated as being between 71 and 475 per 100,000 people as at 11th October.
- The most recent wave of the Scottish Contact Survey (13th 19th October) indicate an average of 5.0 contacts. This has decreased by 18% compared to the previous wave of the survey (29th September -5th October) where average contacts were 6.0. Prior to the Covid-19 pandemic, the average daily contacts for adults in the UK were reported to be 10.8.
- Mean contacts in the other setting (contacts outside home, school and work) have decreased by 40% and in the work setting by 7%. Mean contacts in the home setting have remained at a similar level compared to the previous wave of the survey.
- Mean contacts have decreased in all age groups by at least 6%.

- The largest decrease in interactions compared to the previous wave of the survey was between the 18-29 age group with themselves .
- The largest increases in the proportion of participants visiting different locations is seen in those visiting work (increased from 34% to 38%), healthcare (24% to 28%) and gym settings (19% to 23%).
- The percentage of people wearing a face covering where they have at least one contact outside of the home has decreased from 25% to 22% since the previous wave of the survey pertaining to 29th September – 5th October.
- The percentage of people had taken at least one lateral flow test in the previous 7 days was 26%, the same as in the previous wave of the survey pertaining to 29th September 5th October.
- The SPI-M-O consensus view is that by 22nd November, daily hospitalisations from Covid-19 in Scotland are estimated to be between 7 and 157, and hospital occupancy is estimated to be between 194 and 1609.
- Analysis by the Edinburgh University Roslin Institute indicates that there remains a substantial variation in geographical risk of infection when considering the average recorded census age and booster vaccination uptake.
- The data show a continued decline in reported tests across all deprivation deciles, and a recent rise in LFD positivity. There is no clear trend in positivity with respect to deprivation. The volume of testing remains very low.
- The distribution of lateral flow/LFD tests being reported continues to vary substantially by both age and deprivation status.
- Fewer tests are reported in younger adults across all deciles of deprivation, and for children in more deprived deciles.
- LFD positivity has sharply risen since about 12th September. However, this rise has plateaued in less deprived areas, but continues to rise in more deprived areas. With no deprivation-specific changes in testing volume overall, this is suggestive of disproportionately increased circulation of infection within more deprived communities.
- Nationwide, wastewater Covid-19 levels were between 47 and 89 million gene copies per person per day (Mgc/p/d) during the period 6th 20th October, comparable to the previous two-week period (23rd September 6th October) of 62 to 83 Mgc/p/d.

Overview of Scottish Government Modelling

Modelling outputs are provided here on the current epidemic in Scotland as a whole, based on a range of methods. Because it takes a little over three weeks on average for a person who catches SARS-CoV-2 (the causative agent of Covid-19) to show symptoms, become sick and either die or recover, there is a time lag in what our model can tell us about any changes in the epidemic.

The Scottish Government presents its modelling outputs to the Epidemiology Modelling Review Group (EMRG). These outputs are included (shown in green) in Figure 1. Outputs from other modellers are shown in either cyan (for models based only on cases data) or purple (for models based on other data). The UKHSA consensus range is the rightmost (in red).

The R value and growth rates are also estimated by several independent modelling groups based in universities and the UKHSA. Estimates are considered, discussed and combined at the EMRG, which sits within UKHSA. These are based on data up to 24th October.

The consensus view of the UKHSA across these methods was that the value of R in Scotland is between 0.8 and 1.1 as of 11th October 2022 (Figure 1). Both the lower limit and the upper limit have decreased since the last publication. R is an indicator that lags by two to three weeks.



Figure 1: Estimates of Rt for Scotland as of 11th October, including 90% confidence intervals, produced by EMRG

Source: EMRG

The consensus from UKHSA is that the growth rate in Scotland is between -4% and 3% per day as at 11th October. The lower limit has decreased since the last publication while the upper limit is unchanged.

The various groups which report to EMRG use different sources of data in their models to produce estimates of incidence (Figure 2). The consensus view of the UKHSA across these methods was that the incidence of new daily infections in Scotland was between 71 and 475 new infections per 100,000 people as of 11th October. This equates to between 3,900 and 26,000 people in Scotland becoming infected each day.





What we know about how people's contact patterns have changed

Scottish Contact Survey contact matrices are now published as <u>open</u> <u>data</u>. The dataset contains mean contacts between age groups for all waves of the survey since its inception in August 2020 and will be updated fortnightly to add data for the latest survey wave.

The most recent wave of the Scottish Contact Survey (13th - 19th October) indicate an average of 5.0 contacts. This has decreased by 18% compared to the previous wave of the survey (29th September - 5th October) where average contacts were 6.0, as seen in Figure 3.

Prior to the Covid-19 pandemic, the average daily contacts for adults in the UK were reported to be 10.8 from the UK-wide POLYMOD study¹.

Mean contacts in the other setting (contacts outside home, school and work) have decreased by 40% and in the work setting by 7%. Mean contacts in the home setting have remained at a similar level compared to the previous wave of the survey.

Figure 4 shows how contacts change across age group and setting. Mean contacts have decreased in all age groups by at least 6%.

¹ <u>https://bmcmedicine.biomedcentral.com/counter/pdf/10.1186/s12916-020-01597-8.pdf</u>

Figure 4: Mean contacts per day for adults in Scotland (truncated to 100 contacts per participant) by age group and setting

The heatmaps in Figure 5 show the mean overall contacts between age groups for the surveys relating to 29th September - 5th October and 13th - 19th October and the difference between these periods. The largest decrease in interactions compared to the previous wave of the survey was between the 18-29 age group with themselves.

Figure 5: Mean contacts per day between age groups (truncated to 100 contacts per participant)

The largest increases in the proportion of participants visiting different locations is seen in those visiting work (increased from 34% to 38%), healthcare (24% to 28%) and gym settings (19% to 23%).

Figure 6: Locations visited by participants at least once

Figure 7 shows the percentage of people wearing a face covering where they have at least one contact outside of the home. This has decreased from 25% to 22% since the last wave of the survey .

Figure 7: Percentage of participants wearing a face covering (with at least one contact outside of the home)

In the survey pertaining to 13th - 19th October, 26% of people had taken at least one lateral flow test in the previous 7 days, remaining the same

as in the previous wave of the survey pertaining to the 29th September - 5th October.

Figure 8: Percentage of participants who had taken at least one lateral flow test in the last 7 days

What the modelling tells us about projections of hospitalisations and hospital occupancy in the medium term

SPI-M-O produces projections of the epidemic (Figure 9 and Figure 10), combining estimates from several independent models. These projections are not forecasts or predictions. They represent a scenario in which the trajectory of the epidemic continues to follow the trends that were seen in recent data and do not include the effects of any future policy or behavioural changes. The delay between infection, developing symptoms and the need for hospital care means they cannot fully reflect the impact of behaviour changes in the two to three weeks prior to 24th October. The projections include the potential impact of vaccinations over the next few weeks. Modelling groups have used their expert judgement and evidence from UKHSA, Scottish Universities & Public Health Scotland, and other published efficacy studies when making assumptions about vaccine effectiveness.

*Figure 9: SPI-M-O medium-term projection of daily hospitalisations in Scotland, at 50% and 90% credible intervals*²

Figure 10 shows the SPI-M-O consensus on hospital occupancy. Hospital occupancy is determined by the combination of admissions and length of stay, the latter of which is difficult to model with confidence.

² PHS previously published figures for hospital admissions "with" Covid-19; these have recently changed to admissions "because of" Covid-19. As a result actual admissions figures in these charts are not necessarily directly comparable to those published in previous issues. For more details, please see https://publichealthscotland.scot/our-blog/2022/september/why-we-a re-changing-our-covid-19-reporting/.

Figure 10: SPI-M-O medium-term projection of hospital occupancy in Scotland, at 50% and 90% credible intervals

The SPI-M-O consensus view is that by 22nd November, daily hospitalisations from Covid-19 in Scotland are estimated to be between 7 and 157, and hospital occupancy is estimated to be between 194 and 1609.

Summary of spatial analysis of Covid-19 spread in Scotland

Researchers at the Edinburgh Roslin Institute have conducted spatial analysis of Covid-19 spread in Scotland. A summary of findings from data uploaded up to 20th October 2022 is included here.

Rates of LFD testing are stabilising, albeit at very low levels (0-1 LFD tests reported, per week, per 1,000). PCR testing has fallen (from already very low levels) in recent weeks, with positivity rising.

The distribution of lateral flow/LFD tests being reported varies substantially by both age and deprivation status, with many fewer tests reported in younger adults across all deciles of deprivation, and for children in more deprived deciles. The high level of LFD positivity in the latter category is marked, and when compared to the high number of positives amongst the least deprived, suggests that ascertainment may be lower in younger people in deprived areas.

The sharp rise in LFD positivity since 12th Sept 2022, especially in moderate deprivation deciles and amongst working age adults (Figure 11) is suggestive of increased circulation, and continues to rise for

individuals living in the most deprived areas. However, this rise has plateaued in less deprived areas. With no deprivation-specific changes in testing volume overall, this is suggestive of disproportionately increased circulation of infection within more deprived communities

Figure 11: Variation in testing outcomes comparing Lateral Flow and PCR testing considering age and deprivation status of the data zone of record based on data in week ending 15th October 2022.

Figure 12: Variation in testing outcomes comparing Lateral Flow and PCR testing, separated by deprivation

What can analysis of wastewater samples tell us about local outbreaks of Covid-19 infection?

Levels of Covid-19 RNA in wastewater collected at a number of sites around Scotland are adjusted for population and local changes in intake flow rate (or ammonia levels where flow is not available). See the Technical Annex in Issue 34 of Scottish Government Research Findings for the methodology. These reports are based on the most recent data available. Future updates to data may lead to small retrospective changes.

Nationwide, during the period 6th - 20th October, wastewater COVID-19 levels were between 47 and 89 million gene copies per person per day (Mgc/p/d). As illustrated in Figure 13, the wastewater levels do not give a clear trend and appear to have gone up and down over the two-week period, a similar fluctuating pattern to the previous two-week period.

In Figure 13, we show national wastewater Covid-19 levels up to 20th October 2022, with the ONS Coronavirus Infection Survey (CIS) estimates of Covid-19 prevalence up to 10th October 2022 superimposed.

Figure 13: National running average trends in wastewater Covid-19 as of 20th October³

The relative scale used in Figure 13 to depict the CIS results has been updated to correspond to post-July trends in wastewater RNA, after the ONS survey started using a digital data collection method.

Looking to the future

What may happen in the future around SARS-CoV-2 is uncertain and therefore there are a number of possible Covid-19 futures that may occur in the future. For example, the current Omicron wave may dissipate leaving low levels of Covid-19, or a new variant may emerge potentially having vaccine escape or increased severity, or people's behaviours may change. One approach to this uncertainty is to model alternative versions of the future through the development of different Covid-19 scenarios.

³ For this graph, a wastewater RNA average using the last 7 days of data is computed at every sampling date. Prevalence estimates and 95% confidence intervals from the ONS Coronavirus Infection Survey are overlaid, with a scale chosen to approximately match post-July trends in WW Covid-19.

Given what we know about Covid-19 these possible futures range from a world where immunity reduces Covid-19 hospitalisations and ICU to low levels, through to variant world where a variant with immune escape enters Scotland and Covid-19 hospitalisations and ICU could increase. In between these two extremes there could be possible futures where people's behaviour becomes polarised between those who continue with Covid-19 precautions e.g. hand washing etc. and those who do not.

The scenarios we provide in the next section look at what could happen for planning purposes, not to forecast what will happen. The assumptions are based on our most up to date knowledge, but do not include the effect of future changes in treatment of Covid-19 e.g. widespread use of antivirals or changes in behaviour in response to high levels of infections e.g. in variant world. Therefore, in the most extreme scenarios the peak may be lower than suggested if behaviour or restrictions changed.

There is no linear progression between the worlds and all are plausible. Each world inherently contains a different threat level requiring a different approach to management.

Immune World

In this possible future vaccines and natural immunity are effective at keeping Covid-19 at low levels. New variants may emerge in Scotland but for the foreseeable future infections are based around Omicron.

Infections may decrease from current levels over the coming weeks and months to very low levels. Likewise hospital and ICU occupancy may follow this trend relieving the pressure on healthcare services. Issues with new variants are not considered in this world and therefore levels of infections remain low.

Figure 14: Potential infections, hospital occupancy and ICU occupancy trajectory in Immune World

In Immune world Covid-19 in Scotland reduces below epidemic levels, becoming endemic. Cases of Covid-19 therefore spring up only as rare outbreaks which are controlled through public health measures. People's lives return to something close to normality e.g. physical distancing is not needed but people still choose to self-isolate and hygiene is good. As vaccines are effective, take-up of first/second/third doses are good and boosters become part of an annual cycle like flu. The numbers of people who need medical treatment or hospitalisation for Covid-19 remain low.

The focus moves away from Covid-19 response and into recovery. This includes addressing learning losses, treating Long Covid and working through the hospital backlog. Wellbeing measures improve with reduced anxiety and increased happiness. Those from the highest risk groups feel they can reintegrate without government interventions. The economy continues to recover from the effects of Covid-19. Travellers do not face significant issues with trips overseas.

Polarised world

In this world, vaccines and natural immunity are effective at reducing infections. The approach followed relies on individual risk assessment and behaviours. However, society becomes polarised as some continue to take up vaccines and follow guidance while others are more reluctant. Covid-19 becomes a disease associated with those who do not or cannot get full vaccine benefit and do not or cannot adopt a risk based approach.

Impacts on hospital/ICU occupancy are uncertain but levels may be higher than what may happen in Immune world (see Figure 15).

Cases of Covid-19 spring up and are hard to control in those who are not vaccinated or vulnerable. People's lives return to a "new normal" but,

due to polarised groups in society with some following and some not following guidance, infections remain.

Vaccines are effective so older and more vulnerable people come forward for future doses in high numbers.

The focus remains on Covid-19 and the shift onto recovery is slower. Existing learning losses are harder to rectify and continue to accrue due to infections within education settings. The hospital backlog is difficult to address as hospitals are still dealing with Covid-19 cases. The population becomes polarised in to those whose wellbeing improves e.g. lower risk people and those whose wellbeing deteriorates e.g. higher risk or poorer people whose levels of anxiety increase as Covid-19 circulates. They continue to experience greater illness, greater poverty or disruption to their income. The economy continues to be impacted from the effects of Covid-19.

Variant world – vaccine escape with same severity as Delta

In this possible future a variant with vaccine escape emerges in Scotland presenting a challenge even for fully vaccinated people. This new variant leads to increased transmission, but not to increased severity compared to previous variants. In this scenario other NPIs may need to be put in place for a short time. This world is similar to what has happened in Scotland with the emergence of Omicron.

Omicron may be reduced to low levels within Scotland as a new variant takes over. This causes a new wave of Covid-19 infections as well as increases in hospital and ICU occupancy. People's lives are disrupted due to the increasingly high levels of infections leading to time off work ill or isolating.

To show the potential impact assume a new variant appears in Scotland in late Autumn as was the case with Omicron. The timing is uncertain and a potential new variant may appear sooner than this or significantly later but has currently been lined up in late Autumn to show illustratively what could happen. The new variant may cause Omicron infections to decrease significantly or disappear entirely (and is not shown). The new variant is modelled with similar transmissibility and vaccine escape as Omicron with severity characteristics similar to Delta. It could lead to high levels of infections leading to hospital occupancy rising above capacity restrictions. With sustained high levels of infection we could again see increased staff absences in a number of sectors that were affected by this in the recent Omicron wave.

Figure 16: Potential infections, hospital occupancy and ICU occupancy trajectory in Variant World with the same severity as Delta

The focus remains on Covid-19 and it is hard to shift on to recovery. Continued infections within education settings and staff shortages may impact schools. The Covid-19 strain on hospitals is high due to the very high numbers of infections and workforce pressures grow making it difficult to address the hospital backlog. Wellbeing measures deteriorate with people reporting low happiness and general 'tiredness with it all'. The economy continues to be impacted from the effects of Covid-19 with many people off work. Travellers may not want to come to the UK as the new variant sweeps through.

Variant world – vaccine escape with increased severity compared to Delta

As with the other example of Variant world, a new variant appears in Scotland in late Autumn as was the case with Omicron. The timing is uncertain and a potential new variant may appear sooner than this or significantly later but has currently been lined up in late Autumn to show illustratively what could happen.

The new variant may cause Omicron infections to decrease significantly or disappear entirely (and this is not shown on the graph). It is modelled with similar transmissibility and vaccine escape as Omicron with severity characteristics 50% higher than Delta, purely for illustrative purposes.

It could lead to high levels of infections leading to hospital occupancy rising significantly. With sustained high levels of infection we could again see increased staff absences in a number of sectors that were affected by this in the recent Omicron wave.

Figure 17: Potential infections, hospital occupancy and ICU occupancy trajectory in Variant World with the increased severity compared to Delta

The focus remains on Covid-19 and it is hard to shift on to recovery. Continued infections within education settings and staff shortages may impact schools. The Covid-19 strain on hospitals is high due to the very high numbers of infections and workforce pressures grow making it difficult to address the hospital backlog. Wellbeing measures deteriorate with people reporting low happiness and general 'tiredness with it all'. The economy continues to be impacted from the effects of Covid-19 with many people off work. Travellers may not want to come to the UK as the new variant sweeps through.

What next?

Archiving of models is currently being undertaken via the <u>Data Science</u> <u>Scotland</u> GitHub organisation. Details of the <u>Scottish Contact Survey</u> <u>contact matrices</u> are available. See the Technical Annex of issue 96 for details of other archived models.

Technical Annex

What levels of Covid-19 are indicated by wastewater data?

Table 1 provides population weighted daily averages for normalised wastewater Covid-19 levels in the weeks ending 13th October and 20th October 2022, with no estimate for error. This is given in Million gene copies per person per day. Coverage is given as percentage of inhabitants in each local authority covered by a wastewater Covid-19 sampling site delivering data during this period.

	w/e 13th	w/e 20th	_
Local Authority (LA)	October	October	Coverage
Aberdeen City	135	82	99 %
Aberdeenshire	76	46	44 %
Angus	114	50	68 %
Argyll and Bute	34	60	23 %
City of Edinburgh	134	66	98 %
Clackmannanshire	128	58	92 %
Dumfries and Galloway	78	41	39 %
Dundee City	152	65	100 %
East Ayrshire	122	42	72 %
East Dunbartonshire	119	51	99 %
East Lothian	122	55	74 %
East Renfrewshire	25	31	95 %
Falkirk	93	49	88 %
Fife	67	42	75 %
Glasgow City	59	39	98 %
Highland	70	17	20 %
Inverclyde	12	47	98 %
Midlothian	143	61	88 %
Moray	56	45	70 %
Na h-Eileanan Siar	14	-	0 %
North Ayrshire	81	47	92 %
North Lanarkshire	87	40	80 %
Orkney Islands	32	54	34 %
Perth and Kinross	98	23	45 %
Renfrewshire	42	40	97 %
Scottish Borders	109	73	59 %
Shetland Islands	4	24	29 %
South Ayrshire	127	50	88 %
South Lanarkshire	71	36	85 %
Stirling	16	22	63 %

Table 1. Average Covid-19 wastewater levels (Mgc/p/d)⁴.

⁴ Coverage as for week ending 20th October 2022.

West Dunbartonshire	61	30	98 %
West Lothian	42	40	93 %

This publication will be available in accessible HTML on the gov.scot website

© Crown copyright 2022

You may re-use this information (excluding logos and images) free of charge in any format or medium, under the terms of the Open Government Licence. To view this licence, visit http://www.nationalarchives.gov.uk/doc/opengovernment-licence/ or e-mail: psi@nationalarchives.gsi.gov.uk. Where we have identified any third party copyright information you will need to obtain permission from the copyright holders concerned.

The views expressed in this report are those of the researchers and do not necessarily represent those of the Scottish Government or Scottish Ministers.

This document is also available from our website at www.gov.scot. ISBN: 978-1-80525-156-9

The Scottish Government St Andrew's House Edinburgh EH1 3DG

Produced for the Scottish Government by APS Group Scotland PPDAS1184462 (10/22) Published by the Scottish Government, October 2022

ISBN 978-1-80525-156-9

Web Publication

PPDAS1184462 (10/22)